

REVIEW

The impact of drug interactions and polypharmacy on antimicrobial therapy in the elderly

A. Corsonello¹, A. M. Abbatecola², S. Fusco³, F. Luciani⁴, A. Marino⁵, S. Catalano⁵, M. G. Maggio⁶ and F. Lattanzio²

1) Unit of Geriatric Pharmacoeconomics, Research Hospital of Cosenza, Italian National Research Centre on Aging (INRCA), Cosenza, 2) Scientific Direction, Italian National Research Centre on Aging (INRCA), Ancona, 3) Department of Internal Medicine, University of Messina, Messina, 4) Infectious Diseases Unit, "Annunziata" Hospital, Cosenza, 5) Department of Pharmacy, Health and Nutritional Sciences, and 6) Department of Clinical and Experimental Medicine, Section of Geriatrics, University of Parma, Parma, Italy

Abstract

Infectious diseases are more prevalent in older people than in younger adults, and represent a major healthcare issue in older populations. Indeed, infections in the elderly are often associated with higher morbidity and mortality, and may present atypically. Additionally, older patients are generally treated with polypharmacy regimens, which increase the likelihood of drug–drug interactions when the prescription of an antimicrobial agent is needed. A progressive impairment in the functional reserve of multiple organs may affect either pharmacokinetics or pharmacodynamics during aging. Changes in body composition occurring with advancing age, reduced liver mass and perfusion, and reduced renal excretion may affect either pharmacokinetics or pharmacodynamics. These issues need to be taken into account when prescribing antimicrobial agents to older complex patients taking multiple drugs. Interventions aimed at improving the appropriateness and safety of antimicrobial prescriptions have been proposed. Educational interventions targeting physicians may improve antimicrobial prescriptions. Antimicrobial stewardship programmes have been found to reduce the length of hospital stay and improve safety in hospitalized patients, and their use in long-term care facilities is worth testing. Computerized prescription and decision support systems, as well as interventions aimed at improving antimicrobial agents dosage in relation to kidney function, may also help to reduce the burden of interactions and inherent costs. Clinical Microbiology and Infection © 2014 European Society of Clinical Microbiology and Infectious Diseases. Published by Elsevier Ltd. All rights reserved.

Keywords: Antimicrobial, drug interactions, elderly, pharmacodynamic, pharmacokinetic, polypharmacy

Article published online: 13 October 2014

Corresponding author: A. Corsonello, Unit of Geriatric Pharmacoeconomics, Research Hospital of Cosenza, Italian National Research Centre on Aging (INRCA), C.da Muio Piccolo I-87100 Cosenza, Italy
E-mail: andrea_corsonello@tin.it

Introduction

The choice of the class of antimicrobial agents in older patients is often a challenging issue. Indeed, the type of antibiotic should take into account factors related to patient, culprit pathogen (if and when possible), pharmacokinetic and pharmacodynamic properties, as well as the presence of polypharmacy with the inherent risk of adverse drug reactions, or drug–drug or drug–disease interactions.

The aim of this review is to describe the changes in pharmacokinetics and pharmacodynamics occurring during aging, as well as their impact on polypharmacy and drug interactions involving antimicrobial agents. We focus on clinically relevant interactions with non-antimicrobial medications frequently used in older patients. Potentially useful interventions for reducing the risk of drug interactions when prescribing antimicrobials to older patients are also reviewed.

Age-related changes in pharmacokinetics and pharmacodynamics

Aging is generally characterized by changes in all phases of pharmacokinetic processes (Table 1). However, changes in pharmacokinetics may also result from the co-administration of

TABLE 1. Age-related changes in pharmacokinetics relevant to interactions involving antimicrobial agents

Age related changes		Potential impact on interactions
Absorption	Increased gastric pH	Increased risk of drug-induced oesophageal lesions Changes in solubility and chemical stability of drugs Changes in t_{max} and C_{max} Reduced active transport
	Delayed gastric emptying Reduced splanchnic blood flow Decreased absorption surface Decreased gastrointestinal motility	
Distribution	Changes in body composition	Increased volume of distribution for lipo-soluble drugs Reduced volume of distribution for water-soluble drugs Increased toxicity from selected drugs in the presence of severe hypoalbuminaemia Increased bioavailability of drugs displaced from protein-binding sites Inhibition and/or induction of cytochrome P450s in the context of polypharmacy regimens
	Reduced protein-binding sites Changes in blood–brain barrier permeability (conflicting evidence)	
Metabolism	Reduced hepatic blood flow and overall liver mass Less effective first-pass metabolism and phase I metabolism Reduced cytochrome P450 activity (conflicting evidence)	
Excretion	Reduced kidney glomerular filtration rate and tubular secretion	Impaired elimination of water-soluble drugs

selected drugs, as is the case for several antimicrobial agents, leading to clinically relevant interactions.

Absorption

Reduced oesophageal peristalsis and gastric acid secretion [1] usually have only a minor impact on oral antibiotic absorption. However, an excess increase in gastric pH, as can be produced by long-term use of proton pump inhibitors, can alter solubility and chemical stability of β -lactams, macrolides and azoles, reducing their bioavailability [2]. Age-related reduced gastric emptying and peristalsis [3,4], splanchnic blood flow and bowel surface area [5] can reduce the bioavailability of amoxicillin and clavulanic acid when assumed following the meal [6]. Reduced active transport function may also lead to clinically important drug interactions, which are of particular importance in older patients treated with complex polypharmacy regimens [7,8]. Macrolides may increase serum concentrations of calcium channel blockers and sulphonylureas by inhibiting intestinal cytochrome P450 3A4 (CYP3A4), and cause digoxin toxicity by inhibiting intestinal P-gp [9]. The inhibition of intestinal CYP3A4 by macrolides may increase the risk of toxicity from several drugs, including midazolam, cyclosporine, statins, antiarrhythmics, tricyclic antidepressants, antipsychotics and warfarin. Finally, specific antifungal agents, such as itraconazole and caspofungin, can also inhibit CYP3A4 and P-gp [9].

Distribution

Overall, the volume of distribution for lipophilic drugs may increase with a prolonged half-life [10], whereas water-soluble drugs may have a smaller volume of distribution causing a more rapid increase in plasma concentrations [11], and indicating the need for lower initial doses [12]. Age-related changes in plasma protein binding seem to be less important for drug therapy, as steady-state unbound drug concentration often redistributes and remains unaltered [9,13]. More pronounced changes in protein-binding capacity are generally due to disease-related hypoalbuminaemia [14]. A recent retrospective observational

study of 94 older patients with methicillin-resistant *Staphylococcus aureus* hospital-acquired pneumonia showed that patients with severe hypoalbuminaemia had significantly longer vancomycin half-life (33.2 ± 5.4 versus 24.9 ± 1.6 ; $p = 0.049$), greater risk of 28-day mortality in relation to high values of the area under the concentration curve (AUC)/minimum inhibitory concentration (MIC) ($250\text{--}450$ or $>450 \mu\text{g} \times \text{h/mL}$), and more frequent nephrotoxicity (26% versus 8%, $p < 0.001$) compared with patients with mild hypoalbuminaemia [15]. Hence, severe hypoalbuminaemia needs to be addressed when vancomycin is prescribed to older patients.

Protein-binding drug interactions involving antimicrobial agents also deserve mentioning: co-trimoxazole is known to increase serum concentrations of methotrexate and sulphonylureas by displacing them from plasma protein-binding sites, resulting in a clinically relevant increased risk of hypoglycaemia and severe bone marrow depression [9], respectively.

Metabolism

The bioavailability of drugs undergoing extensive first-pass metabolism increases [16,17], while the bioavailability of drugs that need to be activated in the liver is reduced [18]. In elderly patients, the hepatic clearance of drugs undergoing flow-limited metabolism may be reduced up to 40% [19,20]. Hepatic clearance by cytochrome P450 (CYP)-mediated phase I oxidation, reduction and hydrolysis reactions is impaired to a greater extent with respect to clearance mediated by phase II conjugation reactions, mainly because of the reduced hepatic blood flow and overall liver size [19,20]. The effects of aging on the CYP activities are still a matter of debate [21–23]. An age-related 20% reduction in the metabolism of CYP2D6 substrates has been observed [24,25]. Such a finding has not been confirmed for the CYP3A subfamily, which is responsible for the elimination of more than 50% of its substrates [26–28]. Additionally, the activation of some important CYPs, including CYP3A4, CYP2D6 and CYP1A2, does not seem to change with aging [29,30].

Several antimicrobial agents, mainly macrolides, fluoroquinolones and antifungal azole derivatives undergo metabolic processes in the liver, consisting of phase I metabolism (oxidation) followed by phase II (conjugation) to make drugs hydro-soluble leading to their renal excretion [2,31]. Macrolides, with the exception of azithromycin, inhibit the CYP3A4, and the risk of interaction is greatest with orally administered substrates, such as erythromycin, clarithromycin and telithromycin, which can inhibit both intestinal and hepatic CYP3A4 [9,32,33]. Increased concentrations of selected CYP3A4 substrates, such as midazolam, cyclosporine, tacrolimus, lovastatin, simvastatin and calcium-channel blockers, may have clinically relevant harmful effects, including excessive sedation and falls from benzodiazepines, nephrotoxicity from immunosuppressive agents, rhabdomyolysis from statins, and hypotension from antihypertensive drugs [9,34,35]. Erythromycin and, to a lesser extent, clarithromycin and telithromycin may prolong prothrombin time in patients on warfarin [9,32,33], with consequent increased risk of haemorrhagic events. Enhanced toxicity of phenytoin, sulphonylureas and theophylline has been reported with concomitant use of macrolides [9,32,33,36]. Finally, the inhibition of CYP3A4 by macrolides may also cause fatal drug interactions, such as QTc prolongation leading to torsade de pointe and death due to an increase in levels of some anti-arrhythmics, tricyclic antidepressants and antipsychotic agents [9]. Additionally, by inhibiting CYP3A4, macrolides may increase circulating concentrations of donepezil, a cholinesterase inhibitor used in dementia patients, causing an enhancement in vagal neurotransmission and severe alterations in the sinus node and cardiac conduction systems, sinus bradycardia, neurocardiogenic syncope and bradyarrhythmias [37,38]. Although a recent study failed to find a significantly increased risk of cardiac adverse events in older people on donepezil who were concomitantly exposed to clarithromycin, antimicrobial drugs other than clarithromycin should be preferred whenever possible [39]. Azithromycin lacks the appreciable drug–drug interactions seen with other macrolide antibiotics and is considered the safest of all macrolides from a cardiac perspective. However, the recent finding of a small increase in risk of cardiac events in older patients with pneumonia treated with azithromycin deserves further investigation [40].

By inhibiting CYP3A4 and CYP2C9, fluoroquinolones may enhance the toxicity of selected drugs largely used in older patients, such as benzodiazepines, fentanyl, carbamazepine, statins, theophylline, haloperidol and warfarin. On the contrary, rifampin reduces the bioavailability of warfarin, phenytoin, valproic acid, caspofungin, azoles, digoxin, amiodarone, statins, β -blockers and sulphonylureas by inducing CYP2C9, CYP2C19 and CYP3A4 [9].

Excretion

Renal excretion of drugs can significantly change during advancing age, increasing the risk of toxicity from kidney-cleared antimicrobials. Additionally, several renal excretion interactions involve antimicrobial agents. Plasma concentrations of β -lactams can be increased by drugs affecting their renal tubular secretion, such as probenecid, methotrexate, aspirin and indomethacin. In particular, the co-administration of probenecid doubles the AUCs of amoxicillin, ampicillin, ticarcillin and nafcillin, and increases by 55% the AUC of meropenem [9]. Although higher concentrations of these agents may be desirable for the management of meningitis and endocarditis, the use of probenecid to boost β -lactam concentrations should be avoided in older patients, as well as in those with renal dysfunction, or a history of seizure due to the increased risk of antibiotic-induced convulsions [9,41]. Reduced tubular secretion is also the interaction mechanism by which amantadine, digoxin and methotrexate may increase serum concentration and enhance toxicity of co-trimoxazole [9]. Finally, ciprofloxacin may reduce renal excretion and tubular secretion of methotrexate, leading to severe dermatological, bone marrow, hepatic and renal toxicity [9,42].

Pharmacokinetics/pharmacodynamics

With regards to antimicrobial agents, pharmacodynamics reflects the relationship between serum concentration and the extent to which the drug is able to bind or interact with its specific bacterial target causing cell growth inhibition or death, as measured by MIC [43,44]. Overall, antimicrobial agents may have either a concentration-dependent (e.g. aminoglycosides and quinolones) or a time-dependent (e.g. β -lactams, vancomycin and clindamycin) killing activity. The pharmacodynamic response of concentration-dependent antibiotics largely depends on the maximum plasma level reached. Conversely, the killing activity of time-dependent antibiotics depends on the amount of time during which the plasma antibiotic concentration exceeds the MIC for the organism. For this reason, these agents are optimized by providing smaller, frequent doses or continuous infusion [43–45]. The above considerations clearly indicate that it is very difficult to separate pharmacodynamic processes from pharmacokinetics, and that both age-related changes in pharmacokinetics and pharmacokinetic interactions may obviously affect pharmacodynamics. Indeed, the pharmacokinetics/pharmacodynamics ratio predicts the therapeutic response of microorganisms to antimicrobials by correlating free drug (f) exposure (area under the plasma concentration–time curve over 24 h of dosing ($fAUC_{24}$)) to measures of drug potency (MIC). Hence, the pharmacokinetics/pharmacodynamics parameters able to predict clinical outcomes are the fC_{max}/MIC , the $fAUC_{0-24}/MIC$ (i.e. the area under the

inhibitory plasma concentration-time curve [AUC]), and the time above the MIC (T/MIC) [43]. Using pharmacokinetics/pharmacodynamics parameters for dosing antimicrobials may be useful to achieve therapeutic goals, prevent selection of drug-resistant bacteria and minimize toxic effects when treating infections in older patients [44].

The impact of polypharmacy and interventions to reduce risks

Older patients with co-morbid conditions are frequently excluded from clinical trials [46], and evidence coming from these studies is only partly applicable to this population. This bias also affects clinical practice guidelines that are based on evidence coming from randomized trials and meta-analyses [47]. Guidelines are generally disease-focused, which raises difficulties when applying them in older patients with co-morbid conditions. Indeed, a guideline-driven therapeutic approach in such patients often results in adverse drug–drug or drug–disease interactions in the presence of complex polypharmacy regimens [48].

The risk of potentially dangerous pharmacological interactions involving antimicrobial agents is among the most frequent and undesirable consequences of polypharmacy in older persons. The use of antimicrobial agents in a complex multi-drug regimens was found to be a greater independent risk factor for adverse drug events compared with other drug classes, including antipsychotics or antidepressants, in older nursing home residents [49]. For this reason, lists of clinically important drug interactions, some of which are specific to the elderly [50,51], and recommendations have been proposed [52,53]. Unfortunately, drug interactions are not easily predictable, and nor are variations in specific drug metabolism. It is important to optimize outcomes and to avoid harm from

potentially dangerous interactions by implementing guiding principles for antimicrobial use in older adults. These aims should include the following [52].

- Stratify patient risks for severe infections and multidrug-resistant pathogens based on lifestyle and functional status.
- Provide early empiric therapy using national guidelines and local antibiogram when available.
- Obtain complete medication history and carefully select antimicrobials to avoid interactions.
- Correctly reach maximal therapeutic doses of antimicrobials according to age-related changes in pharmacokinetics and pharmacodynamics to avoid potential adverse effects.
- Discontinue antimicrobial therapy based on the patient's clinical status and identified pathogen.
- Perform clinical trials testing the use of antimicrobial agents in older populations.

Interventions aimed at improving the appropriateness and safety of antimicrobial prescriptions have been investigated (Table 2) [54–61].

Educational interventions targeting prescribing physicians were found to improve antibiotic prescribing practice and to significantly reduce consumption and costs of antibiotics in a geriatric hospital [54].

Antimicrobial stewardship programmes (ASPs), defined as the coordinated effort to optimize antimicrobial usage (the right agent, at the right time, at the correct dose, for an appropriate duration) with the goals of improving patient outcomes, reducing antimicrobial resistance and decreasing healthcare costs [61], have been found to reduce the length of hospital stay and improve safety in hospitalized patients [55]. ASP interventions have also been investigated in long-term care facilities, where residents are often colonized with multidrug-resistant organisms and treated for inappropriately long

TABLE 2. Summary of evidence about interventions specifically aimed at improving appropriateness and safety of antimicrobial prescriptions

Study	Setting and intervention	Evidence
Lutters et al. [54]	Geriatric hospital	Reduced consumption and costs of antibiotics
Liew et al. [55]	Educational intervention targeting prescribing physicians	Reduce length of hospital stay and improved safety
Gonzales et al. [56]	Hospital	
	Antimicrobial stewardship programs	
	Medicare office visits	Modest decline in antibiotic use for acute respiratory infections, but no substantial effect
Bedouch et al. [57]	Educational intervention targeting patients and caregivers	
	Hospital, medical wards	Routine participation of clinical pharmacists in clinical medical rounds may facilitate identification of drug-related problems and enhance patient safety
Rivkin et al. [58]	Computerized physician order entry system and pharmacists	Decreased number of clinically important interactions requiring therapy modification, and reduced length of stay
Buising et al. [59]	Intensive care unit	Improved antibiotic prescribing practices
	Emergency department	
	Computerized decision support system	
Joosten et al. [60]	Ambulatory care setting	
	Automatic renal function alerts (involving general practitioners and community pharmacists)	A considerable proportion of the population is at risk for adverse drug events from antimicrobials due to impaired renal function
		Providing pharmacists and physicians with renal function data may help them to adjust medication dosage

periods, and ASP may be crucial to limit the complications, costs and resistance associated with antibiotic overuse [62]. Although the efficacy of ASPs and how they fit into individual facility types with differing resources still need to be investigated [63], a 30% decrease in systemic antibiotic usage, as well as a decrease in the rate of *C. difficile* infections, has been observed with ASP in a preliminary single-centre study of long-term care facility residents [64].

Computerized support systems have been proved to be effective in identifying antimicrobial-related problems, and increasing the rate of appropriate prescriptions when a clinical pharmacist is involved [57,58]. A computerized decision support system improved the rate of appropriate empirical antibiotic treatment while reducing antibiotic costs and the use of broad-spectrum antibiotic treatment in the context of a cluster randomized controlled trial [65]. Additionally, computerized support system interventions led to a significant improvement in antibiotic prescribing practices in patients presenting to the emergency department with community-acquired pneumonia, and appropriateness of prescriptions was greater than that obtained with academic detailing [59]. Nevertheless, barriers to electronic antibiotic prescribing are still a relevant concern to be addressed to leverage the potential that computerized decision-support systems offer in reducing costs, improving quality and improving patient safety [66].

Interventions to improve antimicrobial selection and dosing in relation to kidney function deserve particular attention. Despite the importance of dosage adjustment among patients with renal impairment, such adjustments are rarely made [67,68]. Indeed, non-compliance with dosing guidelines in hospitalized patients with chronic kidney disease ranges from 19% to 67% [69]. Evidence suggests that reporting estimated glomerular filtration rate (eGFR) may not significantly change prescription habits for nephrotoxic medications, including antibiotics [70,71]. Recently, Joosten et al. [60] showed that introducing automatic renal function alerts in the ambulatory care setting, with the involvement of both general practitioners and community pharmacists, was able to reveal that a considerable proportion of the population is at risk for adverse drug events due to impaired renal function. This type of study provided renal function data to the pharmacists to indicate medication dosage adjustments with prescribing physicians [60]. Hence, it is likely that strategies to implement dosage adjustment in relation to renal function could benefit from the involvement of other professionals involved in the care process, as well as from technology. Indeed, computerized software programs for medication adjustments may improve the appropriateness of prescribed medications and doses [68].

Conclusions

Multiple chronic diseases and complex polypharmacy regimens makes the choice of the right antimicrobial agent very challenging when an infection occurs in older patients. Age-related pharmacokinetic changes play a key role in the risk of drug interactions in these patients [72]. Selected interventions, such as educational interventions, antimicrobial stewardship, computerized prescription and decision support systems, and interventions aimed at improving drug dosage in relation to kidney function, proved to be effective in improving appropriateness and safety when prescribing antimicrobial agents. Hence, the ability of physicians to prescribe safer and more effective antimicrobial therapies when visiting older people with multiple chronic diseases depends substantially on how they will be able to incorporate the principles of drug pharmacokinetics and pharmacodynamics into daily clinical practice [73], as well as how interventions specifically aimed at improving the quality of antimicrobial prescriptions will be implemented in clinical practice.

Transparency Declaration

Conflict of interest disclosure: The authors declare that they have no conflicts of interest.

References

- [1] Feldman M. The mature stomach. Still pumping out acid? *JAMA* 1997;278:681–2.
- [2] Mazzei T. The difficulties of polytherapy: examples from antimicrobial chemotherapy. *Intern Emerg Med* 2011;6(Suppl. 1):103–9.
- [3] Wiley JW. Aging and neural control of the GI tract: III. Senescent enteric nervous system: lessons from extraintestinal sites and nonmammalian species. *Am J Physiol Gastrointest Liver Physiol* 2002;283:G1020–6.
- [4] Orr WC, Chen CL. Aging and neural control of the GI tract: IV. Clinical and physiological aspects of gastrointestinal motility and aging. *Am J Physiol Gastrointest Liver Physiol* 2002;283:G1226–31.
- [5] Ciccocioppo R, Di Sabatino A, Luinetti O, Rossi M, Cifone MG, Corazza GR. Small bowel enterocyte apoptosis and proliferation are increased in the elderly. *Gerontology* 2002;48:204–8.
- [6] Weitschies W, Friedrich C, Wedemeyer RS, Schmidtman M, Kosch O, Kinzig M, et al. Bioavailability of amoxicillin and clavulanic acid from extended release tablets depends on intragastric tablet deposition and gastric emptying. *Eur J Pharm Biopharm* 2008;70:641–8.
- [7] Saltzman JR, Kowdley KV, Perrone G, Russell RM. Changes in small-intestine permeability with aging. *J Am Geriatr Soc* 1995;43:160–4.
- [8] Kinirons MT, Crome P. Clinical pharmacokinetic considerations in the elderly. An update. *Clin Pharmacokinet* 1997;33:302–12.
- [9] Pereira JM, Paiva JA. Antimicrobial drug interactions in the critically ill patients. *Curr Clin Pharmacol* 2013;8:25–38.
- [10] Klotz U, Avant GR, Hoyumpa A, Schenker S, Wilkinson GR. The effects of age and liver disease on the disposition and elimination of diazepam in adult man. *J Clin Invest* 1975;55:347–59.

- [11] Hanratty CG, McGlinchey P, Johnston GD, Passmore AP. Differential pharmacokinetics of digoxin in elderly patients. *Drugs Aging* 2000;17:353–62.
- [12] Morike K, Schwab M, Klotz U. Use of aminoglycosides in elderly patients. Pharmacokinetic and clinical considerations. *Drugs Aging* 1997;10:259–77.
- [13] Rolan PE. Plasma protein binding displacement interactions—why are they still regarded as clinically important? *Br J Clin Pharmacol* 1994;37:125–8.
- [14] Benet LZ, Hoener BA. Changes in plasma protein binding have little clinical relevance. *Clin Pharmacol Ther* 2002;71:115–21.
- [15] Mizuno T, Mizokami F, Fukami K, Ito K, Shibasaki M, Nagamatsu T, et al. The influence of severe hypoalbuminemia on the half-life of vancomycin in elderly patients with methicillin-resistant staphylococcus aureus hospital-acquired pneumonia. *Clin Interv Aging* 2013;8:1323–8.
- [16] Anantharaju A, Feller A, Chedid A. Aging liver. A review. *Gerontology* 2002;48:343–53.
- [17] Tateishi T, Fujimura A, Shiga T, Ohashi K, Ebihara A. Influence of aging on the oxidative and conjugative metabolism of propranolol. *Int J Clin Pharmacol Res* 1995;15:95–101.
- [18] Davies RO, Gomez HJ, Irvin JD, Walker JF. An overview of the clinical pharmacology of enalapril. *Br J Clin Pharmacol* 1984;18(Suppl. 2):215S–29S.
- [19] Schmucker DL. Liver function and phase I drug metabolism in the elderly: a paradox. *Drugs Aging* 2001;18:837–51.
- [20] Le Couteur DG, McLean AJ. The aging liver. Drug clearance and an oxygen diffusion barrier hypothesis. *Clin Pharmacokinet* 1998;34:359–73.
- [21] Shi S, Morike K, Klotz U. The clinical implications of ageing for rational drug therapy. *Eur J Clin Pharmacol* 2008;64:183–99.
- [22] Wauthier V, Verbeeck RK, Calderon PB. The effect of ageing on cytochrome p450 enzymes: consequences for drug biotransformation in the elderly. *Curr Med Chem* 2007;14:745–57.
- [23] Kinirons MT, O'Mahony MS. Drug metabolism and ageing. *Br J Clin Pharmacol* 2004;57:540–4.
- [24] O'Connell MB, Frye RF, Matzke GR, St Peter JV, Willhite LA, Welch MR, et al. Effect of conjugated equine estrogens on oxidative metabolism in middle-aged and elderly postmenopausal women. *J Clin Pharmacol* 2006;46:1299–307.
- [25] Dorne JL, Walton K, Slob W, Renwick AG. Human variability in polymorphic CYP2d6 metabolism: is the kinetic default uncertainty factor adequate? *Food Chem Toxicol* 2002;40:1633–56.
- [26] Parkinson A, Mudra DR, Johnson C, Dwyer A, Carroll KM. The effects of gender, age, ethnicity, and liver cirrhosis on cytochrome p450 enzyme activity in human liver microsomes and inducibility in cultured human hepatocytes. *Toxicol Appl Pharmacol* 2004;199:193–209.
- [27] Schmucker DL. Aging and the liver: an update. *J Gerontol A Biol Sci Med Sci* 1998;53:B315–20.
- [28] Shimada T, Yamazaki H, Mimura M, Inui Y, Guengerich FP. Interindividual variations in human liver cytochrome p-450 enzymes involved in the oxidation of drugs, carcinogens and toxic chemicals: studies with liver microsomes of 30 Japanese and 30 Caucasians. *J Pharmacol Exp Ther* 1994;270:414–23.
- [29] Dilger K, Hofmann U, Klotz U. Enzyme induction in the elderly: effect of rifampin on the pharmacokinetics and pharmacodynamics of propafenone. *Clin Pharmacol Ther* 2000;67:512–20.
- [30] Platten HP, Schweizer E, Dilger K, Mikus G, Klotz U. Pharmacokinetics and the pharmacodynamic action of midazolam in young and elderly patients undergoing tooth extraction. *Clin Pharmacol Ther* 1998;63:552–60.
- [31] Shakunthala N. New cytochrome p450 mechanisms: implications for understanding molecular basis for drug toxicity at the level of the cytochrome. *Expert Opin Drug Metab Toxicol* 2010;6:1–15.
- [32] Pai MP, Graci DM, Amsden GW. Macrolide drug interactions: an update. *Ann Pharmacother* 2000;34:495–513.
- [33] Shi J, Montay G, Bhargava VO. Clinical pharmacokinetics of telithromycin, the first ketolide antibacterial. *Clin Pharmacokinet* 2005;44:915–34.
- [34] Patel AM, Shariff S, Bailey DG, Juurlink DN, Gandhi S, Mamdani M, et al. Statin toxicity from macrolide antibiotic coprescription: a population-based cohort study. *Ann Intern Med* 2013;158:869–76.
- [35] Gandhi S, Fleet JL, Bailey DG, McArthur E, Wald R, Rehman F, et al. Calcium-channel blocker-clarithromycin drug interactions and acute kidney injury. *JAMA* 2013;310:2544–53.
- [36] Pai MP, Momary KM, Rodvold KA. Antibiotic drug interactions. *Med Clin North Am* 2006;90:1223–55.
- [37] Park-Wyllie LY, Mamdani MM, Li P, Gill SS, Laupacis A, Juurlink DN. Cholinesterase inhibitors and hospitalization for bradycardia: A population-based study. *PLoS Med* 2009;6:e1000157.
- [38] Gill SS, Anderson GM, Fischer HD, Bell CM, Li P, Normand SL, et al. Syncope and its consequences in patients with dementia receiving cholinesterase inhibitors: a population-based cohort study. *Arch Intern Med* 2009;169:867–73.
- [39] Hutson JR, Fischer HD, Wang X, Gruneir A, Daneman N, Gill SS, et al. Use of clarithromycin and adverse cardiovascular events among older patients receiving donepezil: a population-based, nested case-control study. *Drugs Aging* 2012;29:205–11.
- [40] Mortensen EM, Halm EA, Pugh MJ, Copeland LA, Metersky M, Fine MJ, et al. Association of azithromycin with mortality and cardiovascular events among older patients hospitalized with pneumonia. *JAMA* 2014;311:2199–208.
- [41] Wallace KL. Antibiotic-induced convulsions. *Crit Care Clin* 1997;13:741–62.
- [42] Dalle JH, Auvrignon A, Vassal G, Leverger G. Interaction between methotrexate and ciprofloxacin. *J Pediatr Hematol Oncol* 2002;24:321–2.
- [43] Mouton JW, Dudley MN, Cars O, Derendorf H, Drusano GL. Standardization of pharmacokinetic/pharmacodynamic (pk/pd) terminology for anti-infective drugs: an update. *J Antimicrob Chemother* 2005;55:601–7.
- [44] Noreddin AM, El-Khatib W, Haynes V. Optimal dosing design for antibiotic therapy in the elderly: a pharmacokinetic and pharmacodynamic perspective. *Recent Pat Antiinfect Drug Discov* 2008;3:45–52.
- [45] Noreddin AM, Haynes V. Use of pharmacodynamic principles to optimise dosage regimens for antibacterial agents in the elderly. *Drugs Aging* 2007;24:275–92.
- [46] Van Spall HG, Toren A, Kiss A, Fowler RA. Eligibility criteria of randomized controlled trials published in high-impact general medical journals: a systematic sampling review. *JAMA* 2007;297:1233–40.
- [47] Pedone C, Lapane KL. Generalizability of guidelines and physicians' adherence. Case study on the sixth joint national committee's guidelines on hypertension. *BMC Public Health* 2003;3:24.
- [48] Boyd CM, Darer J, Boult C, Fried LP, Boult L, Wu AW. Clinical practice guidelines and quality of care for older patients with multiple comorbid diseases: implications for pay for performance. *JAMA* 2005;294:716–24.
- [49] Field TS, Gurwitz JH, Avorn J, McCormick D, Jain S, Eckler M, et al. Risk factors for adverse drug events among nursing home residents. *Arch Intern Med* 2001;161:1629–34.
- [50] Hanlon JT, Schmadre KE. Drug-drug interactions in older adults: which ones matter? *Am J Geriatr Pharmacother* 2005;3:61–3.
- [51] Zhan C, Correa-de-Araujo R, Bierman AS, et al. Suboptimal prescribing in elderly outpatients: potentially harmful drug–drug and drug–disease combinations. *J Am Geriatr Soc* 2005;53:262–7.
- [52] Herring AR, Williamson JC. Principles of antimicrobial use in older adults. *Clin Geriatr Med* 2007;23:481–97. v.
- [53] High KP, Bradley SF, Gravenstein S, Mehr DR, Quagliarello VJ, Richards C, et al. Clinical practice guideline for the evaluation of fever and infection in older adult residents of long-term care facilities: 2008 update by the Infectious Diseases Society of America. *J Am Geriatr Soc* 2009;57:375–94.

- [54] Lutters M, Harbarth S, Janssens JP, Freudiger H, Herrmann F, Michel JP, et al. Effect of a comprehensive, multidisciplinary, educational program on the use of antibiotics in a geriatric university hospital. *J Am Geriatr Soc* 2004;52:112–6.
- [55] Liew YX, Lee W, Loh JC, Cai Y, Tang SS, Lim CL, et al. Impact of an antimicrobial stewardship programme on patient safety in Singapore general hospital. *Int J Antimicrob Agents* 2012;40:55–60.
- [56] Gonzales R, Sauaia A, Corbett KK, Maselli JH, Erbacher K, Leeman-Castillo BA, et al. Antibiotic treatment of acute respiratory tract infections in the elderly: effect of a multidimensional educational intervention. *J Am Geriatr Soc* 2004;52:39–45.
- [57] Bedouch P, Allenet B, Grass A, Labarere J, Brudieu E, Bosson JL, et al. Drug-related problems in medical wards with a computerized physician order entry system. *J Clin Pharm Ther* 2009;34:187–95.
- [58] Rivkin A, Yin H. Evaluation of the role of the critical care pharmacist in identifying and avoiding or minimizing significant drug-drug interactions in medical intensive care patients. *J Crit Care* 2011;26: 104 e101–6.
- [59] Buising KL, Thursky KA, Black JF, MacGregor L, Street AC, Kennedy MP, et al. Improving antibiotic prescribing for adults with community acquired pneumonia: does a computerised decision support system achieve more than academic detailing alone?—a time series analysis. *BMC Med Inform Decis Mak* 2008;8:35.
- [60] Joosten H, Drion I, Boogerd KJ, van der Pijl EV, Slingerland RJ, Slaets JP, et al. Optimising drug prescribing and dispensing in subjects at risk for drug errors due to renal impairment: improving drug safety in primary healthcare by low EGFR alerts. *BMJ Open* 2013;3.
- [61] Rohde JM, Jacobsen D, Rosenberg DJ. Role of the hospitalist in antimicrobial stewardship: a review of work completed and description of a multisite collaborative. *Clin Ther* 2013;35:751–7.
- [62] Daneman N, Gruneir A, Bronskill SE, Newman A, Fischer HD, Rochon PA, et al. Prolonged antibiotic treatment in long-term care: role of the prescriber. *JAMA Intern Med* 2013;173:673–82.
- [63] Rhee SM, Stone ND. Antimicrobial stewardship in long-term care facilities. *Infect Dis Clin North Am* 2014;28:237–46.
- [64] Jump RL, Olds DM, Seifi N, Kyriotakis G, Jury LA, Peron EP, et al. Effective antimicrobial stewardship in a long-term care facility through an infectious disease consultation service: keeping a lid on antibiotic use. *Infect Control Hosp Epidemiol* 2012;33:1185–92.
- [65] Paul M, Andreassen S, Tacconelli E, Nielsen AD, Almasreh N, Frank U, et al. Improving empirical antibiotic treatment using treat, a computerized decision support system: cluster randomized trial. *J Antimicrob Chemother* 2006;58:1238–45.
- [66] Linder JA, Bates DW, Williams DH, Connolly MA, Middleton B. Acute infections in primary care: accuracy of electronic diagnoses and electronic antibiotic prescribing. *J Am Med Inform Assoc* 2006;13:61–6.
- [67] Yap C, Dunham D, Thompson J, Baker D. Medication dosing errors for patients with renal insufficiency in ambulatory care. *Jt Comm J Qual Patient Saf* 2005;31:514–21.
- [68] Hassan Y, Al-Ramahi R, Abd Aziz N, Ghazali R. Drug use and dosing in chronic kidney disease. *Ann Acad Med Singapore* 2009;38:1095–103.
- [69] Long CL, Raebel MA, Price DW, Magid DJ. Compliance with dosing guidelines in patients with chronic kidney disease. *Ann Pharmacother* 2004;38:853–8.
- [70] Geara AS, Azzi N, Ghimire P, Abdallah M, Siddiqui A, Bassil C, et al. The impact of reporting estimated glomerular filtration rate. *Ren Fail* 2011;33:486–8.
- [71] Farag A, Garg AX, Li L, Jain AK. Dosing errors in prescribed antibiotics for older persons with CKD: a retrospective time series analysis. *Am J Kidney Dis* 2014;63:422–8.
- [72] Corsonello A, Pedone C, Incalzi RA. Age-related pharmacokinetic and pharmacodynamic changes and related risk of adverse drug reactions. *Curr Med Chem* 2010;17:571–84.
- [73] Lattanzio F, Corsonello A. The challenge of prescribing drugs in older people. *Curr Drug Metab* 2011;12:599–600.